

## Tobacco and alcohol co-use among people with HIV (PWH) in Nairobi, Kenya

Jonathan Shuter<sup>1</sup>, Katherine Breetz<sup>2</sup>, Christopher W. Kahler<sup>3</sup>, Sylvia A. Ojoo<sup>4</sup>, Patience Oduor<sup>5</sup>, Maureen Ondire<sup>5</sup>, Linda Khakali<sup>5</sup>, Angela O. Achieng<sup>5</sup>, Tina W. Masai<sup>5</sup>, Wendy Potts<sup>6</sup>, Melanie E. Bennett<sup>6</sup>, Lan Li<sup>6</sup>, Emily Koech<sup>5</sup> and Seth S. Himelhoch<sup>7</sup>

<sup>1</sup> Department of Medicine, Albert Einstein College of Medicine, Bronx, USA

<sup>2</sup> University of Kentucky College of Medicine, Lexington, USA

<sup>3</sup> Department of Behavioral and Social Sciences, Brown University School of Public Health, Providence USA

<sup>4</sup> Center for Global Health Practice and Impact, Department of Medicine, Georgetown University Medical Center, Washington, DC, USA

<sup>5</sup> Center for International Health, Education, and Biosecurity-Kenya, University of Maryland, Baltimore, USA

<sup>6</sup> Department of Psychiatry, University of Maryland, Baltimore, USA

<sup>7</sup> Department of Psychiatry, University of Kentucky, Lexington, USA

### Abstract

**Introduction:** Although the majority of people living with HIV (PWH) in the world reside in sub-Saharan Africa, little is known about the syndemic of tobacco and alcohol use among PWH in that region. The purpose of this study was to describe tobacco and alcohol co-use behaviors among PWH in Nairobi, Kenya.

**Methods:** Between June and November 2021, 50 PWH with concurrent tobacco and alcohol use in Nairobi were assessed with a structured interview. Women were intentionally oversampled in order to constitute 50% of the group. Participants completed structured interviews administered by trained research assistants using standardized, culturally appropriate, tobacco and alcohol use questionnaires.

**Results:** Fifty PWH completed the study. All identified as African with a mean age of 40.3 years. All participants were engaged in HIV care and were receiving antiretroviral therapy (ART). Mean cigarettes smoked per day was 8.2, and 57% reported moderate/high cigarette dependence. The mean alcohol use disorder identification test (AUDIT) score was 18.6, and 84% reported drinking amounts in the hazardous range. Preferred alcoholic beverages were clear spirits (48%), beer (32%), and *changaa* (12%), a traditional home-brewed liquor made from millet, corn, or sorghum. Cigarette dependence was not significantly associated with higher AUDIT score in any of the two linear regression models that were tested. In Model 1 younger age ( $\beta = -0.37, p = 0.002$ ) and higher depression score ( $\beta = 0.49, p = 0.02$ ), and in Model 2, younger age ( $\beta = -0.31, p = 0.01$ ) and higher anxiety score ( $\beta = 1.05, p = 0.003$ ) were significantly associated with higher AUDIT scores.

**Conclusions:** Tobacco and alcohol co-use in PWH in Nairobi, Kenya is common. This syndemic may require culturally appropriate and tailored treatment approaches, especially those targeting younger health strata and those with comorbid mental illness, to improve health outcomes.

### Introduction

The Surgeon General implicated cigarette smoking as a cause of lung cancer in 1964 (Public Health Service, Department of Health, Education, and Welfare, 1964), and the World Health Organization (WHO) classified alcohol as a carcinogen in 1988 (WHO International Agency for

Research on Cancer, 1988). A recent detailed analysis of the Global Burden of Disease, Injuries, and Risk Factors dataset demonstrates the importance of tobacco and alcohol use as the first and second leading causes of cancer globally. Tobacco use currently accounts for 36.3% of all cancer deaths among men and 12.3% among women, and alcohol use, the second most important carcinogen in terms of cancer-related mortality, currently accounts for 6.9% of all

**Correspondence:** Jonathan Shuter, 111 East 210th Street, Schiff 2, Bronx, NY 10467, 718-920-7845, 718-405-0610 (fax), [jonathan.shuter@einsteinmed.edu](mailto:jonathan.shuter@einsteinmed.edu)

**Financial support:** This work was supported by award 1R01CA225419 (S.S.H., PI, J.S. and S.A.O., MPI) from the National Institutes of Health (NIH)/National Cancer Institute (NCI) and by the Einstein-Rockefeller-CUNY Center for AIDS Research (P30-AI124414) which is supported by the following NIH co-funding and participating institutes and centers: NIAID, NCI, NICHD, NHBL, NIDA, NIMH, NIA, FIC and OAR. Dr. Kahler's work was facilitated by the Providence/Boston Center for AIDS Research (P30AI042853) and the Alcohol Research Center on HIV (P01AA019072).

**Declaration of interest:** Dr. Shuter has received financial compensation for the development of educational materials and delivering oral presentations from the Northeast/Caribbean AIDS Education and Training Center and from the ObG Project. None of the other authors have competing interests to report.

**Keywords:** HIV, tobacco, smoking, alcohol

cancer deaths among men and 2.3% among women (GBD 2019 Cancer Risk Factors Collaborators, 2022; Sarfati & Gurney, 2022). Tobacco and alcohol are frequently interrelated, as cigarette smokers, on average, drink more and alcohol drinkers, on average, smoke more than their non-using counterparts (Falk et al., 2006; Verplaetse & McKee, 2017). Tobacco and alcohol co-use exerts synergistic effects on some cancer incidences, especially head and neck (Mello et al., 2019) and esophageal malignancies (Prabhu et al., 2014). Although research on tobacco and alcohol co-use predominantly originates from the United States and western Europe, the problem is global (GBD 2019 Cancer Risk Factors Collaborators, 2022).

At the time of the last WHO global status report on alcohol and health (2018), per capita consumption of alcohol was stable in the African region as compared to a 20.3% decrement in the European region (WHO, 2018). Current drinkers in Africa consumed an average of 20% more grams of alcohol per day than the international average, and the age-standardized alcohol-attributable burden of disease and injury was higher in Africa than in any other region of the world (WHO, 2018).

There are high rates of tobacco use among people living with HIV (PWH) across the world with an especially troubling trajectory in sub-Saharan Africa (WHO, 2021). Sub-Saharan Africa is also home to 67% of the world's PWH (Moyo et al., 2023). Throughout sub-Saharan Africa, PWH, both men and women, smoke cigarettes at higher rates than people without HIV (Murphy et al., 2019). Kenyan PWH smoke at higher rates than PWH in any other East African country: 26% of men and 1% of women (Murphy et al., 2019). Studies conducted within sub-Saharan Africa have found that both tobacco smoking (95% CI [1.6, 4.3]) and alcohol drinking (95% CI [1.9, 4.4]) are associated with loss of clinical care follow-up and non-adherence to antiretroviral therapy (ART; Kebede et al., 2021).

This growing body of evidence attests to the importance of studying tobacco and alcohol co-use in PWH in sub-Saharan Africa, the region with the largest overlap of HIV infection with these substances in the world. With the goal of better understanding the co-use of tobacco and alcohol in this region, in particular the drinking patterns in PWH who smoke, and the level of association of alcohol use with tobacco and nicotine dependence, we collected detailed information on tobacco and alcohol use behaviors in a group of Kenyan PWH in the course of conducting a tobacco use treatment trial, and we report our findings herein.

## Methods

This was a cross-sectional interview study of PWH who both smoked cigarettes and consumed alcoholic beverages in Nairobi, Kenya. Study candidates were referred from selected HIV clinics, as well as methadone treatment clinics, or self-referred via pre-approved flyers posted in waiting rooms at HIV treatment sites throughout Nairobi. A sample size of 50 participants was selected, and all of these individuals completed structured interviews in English (three participants), Swahili (40 participants), or a combination of the two languages (seven participants) to

learn about beliefs and practices pertaining to alcohol consumption in PWH who smoked cigarettes in Kenya. Women, who smoke at much lower rates than men in Nairobi (Murphy et al., 2019), were intentionally oversampled in order to constitute 50% of the participant group and allow for meaningful descriptive analyses of both sexes. Participants were screened in clinics on scheduled screening visit days with inclusion/exclusion criteria as listed below. These participants were co-enrolled into an ongoing smoking cessation treatment trial (NCT03342027).

**Inclusion criteria:** For inclusion in the study, patients were required to be 18 years or older with a chart diagnosis of HIV, self-reported cigarette smoking of at least one puff in the last 7 days, and any reported alcohol use in the last 30 days. Additionally, participants were required to receive care at one of the assigned recruitment sites and to provide informed consent to participate. **Exclusion criteria:** Potential participants were excluded if study physicians believed they were not medically stable as determined by review of past medical history and current medical status, if they had plans to relocate or travel for more than two consecutive months during the smoking cessation trial, or if their expected survival was less than six months.

Eligible patients were assisted by trained research assistants with consent in writing for participation in either Swahili or English according to the patient's preference.

Table 1 summarizes the tobacco and alcohol use scales that we administered (Abrams & Biener, 1992; Bohn et al., 1995; Fagerstrom & Schneider, 1989; Keyson M, n.d.; Kiluk et al., 2013; Leigh & Stacy, 1993; McKiernan et al., 2011; Saunders et al., 1993). In addition to these, we used individual items from the Centers for Disease Control Question Inventory on Tobacco (Centers for Disease Control and Prevention, 2023) and questions of our own design to collect culturally relevant information from the study sample. The latter questions were reviewed, edited, and approved by native Nairobi residents who were members of the investigative team and included such items as *changaa* and *kumi-kumi* among the choices of alcohol consumed. Most common alcohol used was stratified by first generation (e.g. Smirnoff), second generation (e.g. Kenya Cane), and traditional brew (e.g. Busaa). The Alcohol Use Disorders Identification Test (AUDIT) is a widely endorsed 10-item screening test for alcohol use disorder (AUD; O'Connor et al., 2018), with outcome categories of low risk for AUD (scores = 0–7), medium risk (8–15), high risk (16–19), and addiction likely (20–40). The AUDIT scale has been deployed in Kenyan populations (Luchters et al., 2011), and it has served as an outcome variable in analyses of predictors and correlates of AUD in prior research (Murphy & Garavan, 2011). Other illicit drug use, including marijuana, cocaine, heroin, amphetamines, glue, and prescription drugs, was also assessed, as was methadone treatment status, for both current exposure (i.e. past 30 days) and lifetime exposure. Finally, mental health was addressed with questions pertaining to depression (Center for Epidemiologic Studies Short Depression Scale [CES-D 10]) and anxiety (Generalized Anxiety Disorder 7-item [GAD-7]), both of which have been associated with tobacco and alcohol use behaviors in PWH (Garey et al., 2015; Mdodo et

al., 2015; Zyambo et al., 2015). Quantitative data were collected and stored via Research Electronic Data Capture (REDCap) allowing for encrypted patient information compliant with Health Insurance Portability and

Accountability Act of 1996 guidelines. Participants were compensated with 500 Kenyan Shillings to facilitate travel to complete the study visit.

**Table 1**

***Tobacco and Alcohol Use Measures***

Measure	#Items	Range [Interpretation]	Ref
<b>Tobacco measures</b>			
Fagerstrom test for cigarette dependence	6	0-10 (higher score = greater dependence)	Fagerstrom & Schneider, 1989
Abrams-Biener readiness to quit ladder	1	1-10 (higher score = greater motivation to quit)	Abrams & Beiner, 1992
<b>Alcohol measures</b>			
AUDIT	10	0-40 (higher score = higher risk for alcohol use disorder)	Saunders et al., 1993
Alcohol Outcome Expectancies questionnaire	34	1-6 (Positive subscales: social facilitation/fun/sex/tension reduction. Higher score=greater expectancy. Negative subscales: social/emotional/physical/cognitive performance. Higher score=greater expectancy)	Leigh & Stacy, 1993
Short Inventory of Problems	15	Total score 0-45 All subscales 0-9 (Higher score = more frequent occurrence of problem)	Kiluk et al., 2013
Alcohol Urge Questionnaire	7	7-49 (Higher score = greater urge)	Bohn et al., 1995
Alcohol abstinence self-efficacy	12	1-5 (Total score and subscales: positive affect/negative affect/craving/physical. Higher score = greater self-efficacy)	McKiernan et al., 2011
Drinking-related internal-external locus of control	25	Internal 1-42 External 1-42 Fate 1-21 (Higher score=greater estimation of control)	Keyson, n.d.

Continuous variables were summarized with means and standard deviations. Comparisons of means were accomplished using Student's *t*-test and bivariate correlation of continuous variables was accomplished with Pearson's product moment correlation. In order to evaluate predictors of higher levels of alcohol use and risk for alcohol use disorder, we employed multiple linear regression with AUDIT score as the outcome variable, and all measures having statistically significant associations with AUDIT score on bivariate analyses were included in the model. Statistical analyses were completed using the SAS 9.4M7 software package.

All aspects of the study were approved and overseen by the University of Maryland Institutional Review Board and the Ethics and Research Committee of the Kenyatta National Hospital/University of Nairobi.

## Results

We enrolled 50 participants into this trial of pharmacologic and/or behavioral cessation therapy between June and November 2021. The sociodemographic and clinical characteristics of the cohort and their scores on the various scales relating to cigarette and alcohol use are summarized in Table 2. The mean age of participants was  $40.3 \pm 10.2$  years, 50% were male, and 50% were female. The majority

of participants, 84%, did not complete secondary education, 63% worked full-time outside of the home, and 92% reported stable housing, i.e. living in one's own apartment or house as opposed to being hosted temporarily by family or friends or to being undomiciled. Approximately one-third of participants were married. The majority of participants reported HIV exposure and infection from heterosexual contact, and all reported currently taking ART medications. Fifteen of the participants were receiving methadone maintenance therapy.

Participants on average were  $20.0 \pm 6.8$  years old when they first started smoking tobacco cigarettes, and 87.8% of participants reported they were current, everyday smokers. The average number of cigarettes per day was  $8.2 \pm 6.9$ . On the Fagerstrom Test for Cigarette Dependence (FTCD; Fagerstrom & Schneider, 1989), 14.9% had low dependence, 27.7% had low-moderate dependence, 53.2% had moderate dependence, and 4.3% had high nicotine dependence. Motivation to quit was high with 67.4% in the preparation stage, 30.6% in the contemplation stage, and 2.0% in the precontemplation stage on the Abrams-Biener Readiness to Quit Ladder (Abrams & Biener, 1992).

Participants on average were  $22.2 \pm 6.2$  years old when they first began to drink alcohol. Most commonly participants reported consumption of first (32%) and second (48%) generation alcohol containing beverages (see Table 2 for

definitions of alcohol generation categories), as well as *changaa* (14%), a traditional home-brewed liquor made from millet, corn, or sorghum. Preferred beverages included clear spirits (48%), beer (32%), and *changaa* (12%). On the AUDIT scale, eight (16%) were at low risk (scores = 0–7),

eleven (22%) were at medium risk (8–15), and six (12%) were at high risk (16–19) for alcohol use disorder; and the remaining 25 (50%), the highest score stratum, were classified as “addiction likely” (20–40; WHO, 2001).

**Table 2*****Sociodemographic, Clinical, and Behavioral Characteristics of the Study Sample***

Characteristic or Measure		<i>n</i> (%) or Mean±SD
Age (years)		40.3±10.2
Sex	Male	25 (50%)
	Female	25 (50%)
Ethnicity	Kikuyu	29 (58%)
	Kamba	9 (18%)
	Luo	4 (8%)
	Other or multiple <sup>a</sup>	8 (16%)
Religion	Catholic	25 (50%)
	Protestant	17 (34%)
	Muslim or other <sup>b</sup>	8 (16%)
Marital status	Married/living with partner	15 (30%)
	Single	11 (22%)
	Separated or divorced	20 (40%)
	Widowed	4 (8%)
Housing status	Stable	46 (92%)
	Transitional or homeless	4 (8%)
Employment status	Full-time	14 (30.4%)
	Part-time	29 (63.0%)
	Unemployed	3 (6.5%)
Educational attainment	Completed some primary education	15 (30%)
	Completed primary education	9 (18%)
	Completed some secondary education	15 (30%)
	Completed secondary education ± college education	8 (16%)
	No education	3 (6%)
Mode of HIV acquisition	Heterosexual contact	26 (52%)
	Same-sex contact	4 (8%)
	Injection drug use	9 (18%)
	Other/unknown	11 (22%)
Daily cigarette consumption		8.2±6.9
Fagerstrom test for cigarette dependence		4.5±2.2
Most common type of alcohol consumed	1st generation (most expensive) <sup>d</sup>	16 (32%)
	2nd generation (less expensive) <sup>e</sup>	24 (48%)
	<i>Changaa</i>	7 (14%)
	Other traditional brews <sup>f</sup>	3 (6%)
Preferred category of alcohol consumed	Clear spirits (e.g. rum, gin, tequila)	24 (48%)
	Beer	16 (32%)
	<i>Changaa</i>	6 (12%)
	Whiskey, wine, or champagne	4 (8%)
Most common location of alcohol consumption	Bar or nightclub	30 (60%)
	Home	10 (20%)
	Other <sup>c</sup>	10 (20%)
AUDIT		18.6 ± 9.8
Alcohol outcome expectancies (subscales)	Social facilitation	4.0±1.2
	Improved sexual experience	3.1±1.5
	Fun	4.1±1.1
	Tension reduction	4.1±1.2
Short Inventory of Problems		20.7±13.3
Alcohol Urge Questionnaire		20.6±12.4
Alcohol abstinence self-efficacy		2.7±1.0

Characteristic or Measure		<i>n</i> (%) or Mean±SD
Drinking-related internal-external locus of control (subscales)	Internal	23.1±10.4
	External	28.6±9.2
	Fate	12.1±4.0
Other substance use	Marijuana/bhang/K2/hashish/THC	
	Ever	30 (60%)
	Past 30 days	17 (34%)
	Cocaine <sup>g</sup>	10 (20%)
	Heroin <sup>h</sup>	13 (26%)
	Amphetamines/uppers/ecstasy/miraa/khat	
	Ever	33 (66%)
	Past 30 days	17 (34%)
Sniffed glue	Ever	4 (8%)
	Past 30 days	4 (8%)
CES-D <sup>i</sup>		7.6±6.5
GAD-7 <sup>i</sup>		3.5±3.7

**Notes:**<sup>a</sup> Includes: Luhya (2), Meru (1), Somali (1), Taita (1), other or multiple (3)<sup>b</sup> Includes: Muslim (6), other or no religion (2)<sup>c</sup> Includes: Restaurant (1), work (1), home of friend or relative (1), other (7)<sup>d</sup> E.g. Amarula, Viberov, Smirnoff, Richot, Tusker, Pilsner, Guinness.<sup>e</sup> E.g. Senator Keg, Kenya Cane, Iceberg, Konyagi, Marry Cane<sup>f</sup> E.g. *Pombe za kitamaduni, kama, busaa, muratina*, or other type of alcohol<sup>g</sup> No participants reported cocaine use within the past 30 days.<sup>h</sup> One participant reported heroin use within the past 30 days.<sup>i</sup> CES-D=Center for epidemiologic studies – depression scale; GAD-7=General anxiety disorder – 7 scale.

Bivariate correlations (Pearson's) were completed to evaluate associations of age, nicotine dependence, CES-D and GAD-7 scores, with AUDIT scores. Significant

relationships were discovered between age, depression, anxiety, and AUDIT scores in our sample, as listed in Table 3.

**Table 3*****Bivariate Correlation and Multiple Linear Regressions evaluating Relationships between Variables with AUDIT Score***

Variable	Pearson's <i>r</i>	<i>p</i>
Age (years)	-0.49	<0.001
Cigarette dependence	0.27	0.058
CES-D	0.48	<0.001
GAD-7	0.55	<0.001

**Multiple Linear Regression: Model 1**

Covariate	$\beta^a$ [95% C.I.]	<i>p</i>	<i>R</i> <sup>2</sup>
Age (years)	-0.37 [-0.61, -0.14]	0.002	0.24
Cigarette dependence <sup>b</sup>	0.58 [-0.51, 1.67]	0.29	0.07
CES-D	0.49 [0.10, 0.88]	0.02	0.23

**Multiple Linear Regression: Model 2**

Covariate	$\beta^a$ [95% C.I.]	<i>p</i>	<i>R</i> <sup>2</sup>
Age (years)	-0.31 [-0.55, -0.07]	0.01	0.24
Cigarette dependence <sup>b</sup>	0.77 [-0.24, 1.78]	0.13	0.07
GAD-7	1.05 [0.37, 1.73]	0.003	0.31

**Notes:**

AUDIT=Alcohol use disorders identification test; C.I.=confidence intervals.

Model 1 includes age, cigarette dependence and CES-D score as covariates. Model 2 includes age, nicotine dependence, and GAD-7 score as covariates.

<sup>a</sup> The regression coefficient is calculated for a one-unit change for each covariate.<sup>b</sup> Nicotine dependence is derived from the Fagerstrom test for cigarette dependence (Fagerstrom & Schneider, 1989).

In the multivariate analysis, depression and anxiety scores demonstrated a high degree of collinearity, and thus separate

models were created including depression score (and excluding anxiety score) and including anxiety score (and

excluding depression score). In the adjusted model that included depression score, only age ( $\beta = -0.37$ , 95% CI [-0.61, 0.14],  $p = 0.002$ ) and depression ( $\beta = 0.49$ , 95% CI [-0.10, 0.88],  $p = 0.02$ ) retained their significant associations with AUDIT score (Table 3). In the adjusted model that included anxiety score, only age ( $\beta = -0.31$ , 95% CI [-0.55, 0.07],  $p = 0.01$ ) and anxiety ( $\beta = 1.05$ , 95% CI [0.37, 1.73],  $p = 0.003$ ) retained their significant positive associations with AUDIT score (Table 3).

## Discussion

The literature exploring and describing strategies to treat tobacco use and alcohol use as individual behaviors is voluminous and spans many decades. However, although the interrelationships among various substance uses are extensive, profoundly affecting the outcomes of treatment efforts, the literature on the management of substance co-use, including tobacco and alcohol co-use, is sparse. Although the interactions between these two substances have been recognized since (at least) the early 19<sup>th</sup> century (Fowler, 1842), modern research sheds light on the physiological and sociological interactions between tobacco and alcohol use and acknowledges the obstacle that addiction to one substance presents when attempting to reduce or quit the other one (Frie et al., 2022).

In the present study we describe a sample of PWH from Nairobi, who were both current cigarette smokers and alcohol drinkers. Participants smoked an average of eight cigarettes per day with moderate nicotine dependence, and motivation to quit was high, albeit in a sample of individuals expressing interest in a smoking cessation trial. Our survey of alcohol use patterns revealed that 84% of participants scored in the hazardous drinking range on the AUDIT scale. Generally, participants preferred clear spirits and beer, and second generation (less expensive) beverages were consumed most often.

Younger age, anxiety, and depression were all found to be predictive of higher AUDIT scores. These results are consistent with earlier literature describing research conducted in other regions including one study that correlated moderate/severe anxiety with higher AUDIT scores and older age with lower AUDIT scores and alcohol severity (Silverberg et al., 2018). Another study found similar evidence of higher AUDIT scores associated with anxiety, depression, and other drug use (Khan et al., 2020). Efforts to treat tobacco use in PWH have met with limited success to date, and it is possible that failing to account for co-use of other substances, such as alcohol, and the behavioral and psychological correlates of such co-use may be contributing to this ongoing public health challenge.

Our study has certain limitations deserving of mention. The sample size was comparatively small, and this could have limited our ability to define certain summary variables with precision or to detect differences between subgroups, in particular the association between nicotine dependence and AUDIT score. The paper and pencil interview format conducted in the presence of research staff could have led to social desirability bias in responses. Since participants were being screened for inclusion in a smoking cessation trial, the

sample was preselected for interest in quitting. The characteristics of unmotivated tobacco and alcohol co-users may differ from our sample. Finally, we enrolled in one large urban center in sub-Saharan Africa, and our findings may not be generalizable to other cities, countries, or regions.

The enormous human tragedy caused by the HIV epidemic continues in Sub-Saharan Africa, and the gains achieved by expanded access to antiretroviral therapy are being partially offset by various substance uses, especially tobacco and alcohol. Tobacco and alcohol co-use is common and is associated with younger age and with higher levels of depression and anxiety. Effective treatments for tobacco use and for alcohol use disorder should pay heed to the characteristics of co-users, consider the inclusion of mood management strategies for heavier drinkers, and craft their content to target the appropriate age strata. Public health priorities in much of the world have begun to focus on the burden of non-communicable diseases among PWH, and our research supports the inclusion and implementation of tobacco, alcohol, and substance use treatment together with mental health screening and management as vital components of comprehensive HIV care.

## Acknowledgements

The authors acknowledge with gratitude the cooperation of the study participants in permitting us to complete this research.

## References

- Abrams, D. B., & Biener, L. (1992). Motivational characteristics of smokers at the workplace: A public health challenge. *Preventive Medicine*, 21(6), 679–687. [https://doi.org/https://doi.org/10.1016/0091-7435\(92\)90075-S](https://doi.org/https://doi.org/10.1016/0091-7435(92)90075-S)
- Bohn, M. J., Krahn, D. D., & Staehler, B. A. (1995). Development and initial validation of a measure of drinking urges in abstinent alcoholics. *Alcoholism, Clinical and Experimental Research*, 19(3), 600–606. <https://doi.org/10.1111/j.1530-0277.1995.tb01554.x>
- Centers for Disease Control and Prevention. (2023). Question Inventory on Tobacco (QIT). Retrieved on October 4, 2023, from <https://chronicdata.cdc.gov/Survey-Questions-Tobacco-Use-/Question-Inventory-on-Tobacco-QIT-/vdgb-f9s3/data>
- Fagerstrom, K. O., & Schneider, N. G. (1989). Measuring nicotine dependence: A review of the Fagerstrom Tolerance Questionnaire. *Journal of Behavioral Medicine*, 12(2), 159–182. <https://doi.org/10.1007/BF00846549>
- Falk, D. E., Yi, H. Y., & Hiller-Sturmhofel, S. (2006). An epidemiologic analysis of co-occurring alcohol and tobacco use and disorders: Findings from the National Epidemiologic Survey on Alcohol and Related Conditions. *Alcohol Research & Health*, 29(3), 162–171. <https://www.ncbi.nlm.nih.gov/pubmed/17373404>
- Fowler, O. (1842). *A disquisition on the evils of using tobacco*. Geo. Gregory Publishers,.
- Frie, J. A., Nolan, C. J., Murray, J. E., & Khokhar, J. Y. (2022). Addiction-related outcomes of nicotine and alcohol co-use: New insights following the rise in vaping. *Nicotine & Tobacco Research*, 24(8), 1141–1149. <https://doi.org/10.1093/ntr/ntab231>
- Garey, L., Bakhshaie, J., Sharp, C., Neighbors, C., Zvolensky, M. J., & Gonzalez, A. (2015). Anxiety, depression, and HIV symptoms among persons living with HIV/AIDS: The role of

- hazardous drinking. *AIDS Care*, 27(1), 80–85. <https://doi.org/10.1080/09540121.2014.956042>
- GBD 2019 Cancer Risk Factors Collaborators (2022). The global burden of cancer attributable to risk factors, 2010–19: A systematic analysis for the Global Burden of Disease Study 2019. *Lancet*, 400(10352), 563–591. [https://doi.org/10.1016/S0140-6736\(22\)01438-6](https://doi.org/10.1016/S0140-6736(22)01438-6)
- Kebede, H. K., Mwanri, L., Ward, P., & Gesesew, H. A. (2021). Predictors of lost to follow up from antiretroviral therapy among adults in sub-Saharan Africa: A systematic review and meta-analysis. *Infectious diseases of poverty*, 10(1), 33. <https://doi.org/10.1186/s40249-021-00822-7>
- Keyson M, J. L. (n.d.). Drinking Related Internal-External Locus of Control Scale . Retrieved on February 17, 2023 from [https://www.drugsandalcohol.ie/26771/1/Drinking\\_Related\\_Internal\\_External\\_Locus\\_of\\_Control\\_Scale\\_103.pdf](https://www.drugsandalcohol.ie/26771/1/Drinking_Related_Internal_External_Locus_of_Control_Scale_103.pdf).
- Khan, M. R., Young, K. E., Caniglia, E. C., Fiellin, D. A., Maisto, S. A., Marshall, B. D. L., Edelman, E. J., Gaither, J. R., Chichetto, N. E., Tate, J., Bryant, K. J., Severe, M., Stevens, E. R., Justice, A., & Braithwaite, S. R. (2020). Association of alcohol screening scores with adverse mental health conditions and substance use among US Adults. *JAMA Network Open*, 3(3), e200895. <https://doi.org/10.1001/jamanetworkopen.2020.0895>
- Kiluk, B. D., Dreifuss, J. A., Weiss, R. D., Morgenstern, J., & Carroll, K. M. (2013). The Short Inventory of Problems – Revised (SIP-R): Psychometric properties within a large, diverse sample of substance use disorder treatment seekers. *Psychology of Addictive Behaviors*, 27(1), 307–314. <https://doi.org/10.1037/a0028445>
- Leigh, B. C., & Stacy, A. W. (1993). Alcohol outcome expectancies: Scale construction and predictive utility in higher order confirmatory models. *Psychological Assessment*, 5(2), 216–229. <https://doi.org/10.1037/1040-3590.5.2.216>
- Luchters, S., Geibel, S., Syengo, M., Lango, D., King'ola, N., Temmerman, M., & Chersich, M. F. (2011). Use of AUDIT, and measures of drinking frequency and patterns to detect associations between alcohol and sexual behaviour in male sex workers in Kenya. *BMC Public Health*, 11, 384. <https://doi.org/10.1186/1471-2458-11-384>
- McKiernan, P., Cloud, R., Patterson, D. A., Wolf Adelv Uneyg Waya, S., Golder, S., & Besel, K. (2011). Development of a brief abstinence self-efficacy measure. *Journal of Social Work Practice in the Addictions*, 11(3), 245–253. <https://doi.org/10.1080/1533256X.2011.593445>
- Mdodo, R., Frazier, E. L., Dube, S. R., Mattson, C. L., Sutton, M. Y., Brooks, J. T., & Skarbinski, J. (2015). Cigarette smoking prevalence among adults with HIV compared with the general adult population in the United States: Cross-sectional surveys. *Annals of Internal Medicine*, 162(5), 335–344. <https://doi.org/10.7326/M14-0954>
- Mello, F. W., Melo, G., Pasetto, J. J., Silva, C. A. B., Warnakulasuriya, S., & Rivero, E. R. C. (2019). The synergistic effect of tobacco and alcohol consumption on oral squamous cell carcinoma: A systematic review and meta-analysis. *Clinical Oral Investigations*, 23(7), 2849–2859. <https://doi.org/10.1007/s00784-019-02958-1>
- Moyo, E., Moyo, P., Murewanhema, G., Mhango, M., Chitungo, I., & Dzinamarira, T. (2023). Key populations and Sub-Saharan Africa's HIV response. *Frontiers in Public Health*, 11, 1079990. <https://doi.org/10.3389/fpubh.2023.1079990>
- Murphy, J. D., Liu, B., & Parascandola, M. (2019). Smoking and HIV in sub-Saharan Africa: A 25-country analysis of the Demographic Health Surveys. *Nicotine & Tobacco Research*, 21(8), 1093–1102. <https://doi.org/10.1093/ntr/nty176>
- Murphy, P., & Garavan, H. (2011). Cognitive predictors of problem drinking and AUDIT scores among college students. *Drug and Alcohol Dependence*, 115(1-2), 94–100. <https://doi.org/10.1016/j.drugalcdep.2010.10.011>
- O'Connor, E. A., Perdue, L. A., Senger, C. A., Rushkin, M., Patnode, C. D., Bean, S. I., & Jonas, D. E. (2018). Screening and behavioral counseling interventions to reduce unhealthy alcohol use in adolescents and adults: Updated evidence report and systematic review for the US Preventive Services Task Force. *JAMA*, 320(18), 1910–1928. <https://doi.org/10.1001/jama.2018.12086>
- Prabhu, A., Obi, K. O., & Rubenstein, J. H. (2014). The synergistic effects of alcohol and tobacco consumption on the risk of esophageal squamous cell carcinoma: a meta-analysis. *American Journal of Gastroenterology*, 109(6), 822–827. <https://doi.org/10.1038/ajg.2014.71>
- Sarfati, D., & Gurney, J. (2022). Preventing cancer: The only way forward. *Lancet*, 400(10352), 540–541. [https://doi.org/10.1016/S0140-6736\(22\)01430-1](https://doi.org/10.1016/S0140-6736(22)01430-1)
- Saunders, J. B., Aasland, O. G., Babor, T. F., De La Fuente, J. R., & Grant, M. (1993). Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO Collaborative Project on Early Detection of Persons with Harmful Alcohol Consumption-II. *Addiction*, 88(6), 791–804. <https://doi.org/https://doi.org/10.1111/j.1360-0443.1993.tb02093.x>
- Silverberg, M. J., Leyden, W. A., Leibowitz, A., Hare, C. B., Jang, H. J., Sterling, S., Catz, S. L., Parthasarathy, S., Horberg, M. A., & Satre, D. D. (2018). Factors associated with hazardous alcohol use and motivation to reduce drinking among HIV primary care patients: Baseline findings from the Health & Motivation study. *Addictive Behaviors*, 84, 110–117. <https://doi.org/10.1016/j.addbeh.2018.03.033>
- Public Health Service, Department of Health, Education, and Welfare. (1964, January 1). Smoking and Health: Report of the Advisory Committee to the Surgeon General of the Public Health Service. [Government]. Retrieved on December 6, 2022, from <https://www.govinfo.gov/app/details/GPO-SMOKINGANDHEALTH>
- Verplaetse, T. L., & McKee, S. A. (2017). An overview of alcohol and tobacco/nicotine interactions in the human laboratory. *American Journal of Drug and Alcohol Abuse*, 43(2), 186–196. <https://doi.org/10.1080/00952990.2016.1189927>
- World Health Organization. (2018). *Global status report on alcohol and health 2018*. World Health Organization. Licence: CC BY-NC-SA 3.0 IGO.
- World Health Organization. (2021). *Status of tobacco production and trade in Africa*. World Health Organization. Licence: CC BY-NC-SA 3.0 IGO.
- World Health Organization, Babor, Thomas F., Higgins-Biddle, John C., Saunders, John B. & Monteiro, Maristela G. (2001). *AUDIT: The alcohol use disorders identification test: Guidelines for use in primary health care*, 2nd ed. World Health Organization. Retrieved on February 22, 2023, from <https://iris.who.int/handle/10665/67205>
- World Health Organization. International Agency for Research on Cancer. (1988). Alcohol drinking. *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*. 44, 1–378.
- Zyambo, C. M., Willig, J. H., Cropsey, K. L., Carson, A. P., Wilson, C., Tamhane, A. R., Westfall, A. O., & Burkholder, G. A. (2015). Factors associated with smoking status among HIV-positive patients in routine clinical care. *Journal of AIDS & Clinical Research*, 6(7). <https://doi.org/10.4172/2155-6113.1000480>